

L^2 -Solutions for Nonlinear Schrödinger Equations and Nonlinear Groups

By

Yoshio TSUTSUMI

(Hiroshima University, Japan)

§1. Introduction and main results

We consider the unique global existence of solutions in a weaker class than $H^1(\mathbf{R}^n)$ and some properties of the solution operator for the following nonlinear Schrödinger equation:

$$(1.1) \quad i \frac{\partial u}{\partial t} = -\Delta u + \lambda |u|^{p-1} u, \quad t \in \mathbf{R}, \quad x \in \mathbf{R}^n,$$

$$(1.2) \quad u(t_0, x) = u_0(x),$$

where $t_0 \in \mathbf{R}$ and $\lambda \in \mathbf{R}$. By $\alpha(n)$ we denote ∞ if $n=1$ or $n=2$ and $(n+2)/(n-2)$ if $n \geq 3$. There are many papers concerning the global existence of solutions for Problem (1.1)–(1.2) (see, e.g., [1]–[2], [4]–[7] and [9]). In [1] Baillon, Cazenave and Figueira show that if $1 \leq n \leq 3$, $1 < p < \alpha(n)$ and $\lambda > 0$, Problem (1.1)–(1.2) has a unique global strong solution $u(t) \in C(\mathbf{R}; H^2(\mathbf{R}^n)) \cap C^1(\mathbf{R}; L^2(\mathbf{R}^n))$ for any $u_0 \in H^2(\mathbf{R}^n)$. In [2] Ginibre and Velo show that if $1 < p < \alpha(n)$ and $\lambda > 0$ or if $1 < p < 1 + 4/n$ and $\lambda < 0$, Problem (1.1)–(1.2) has a unique global weak solution $u(t) \in C(\mathbf{R}; H^1(\mathbf{R}^n))$ for any $u_0 \in H^1(\mathbf{R}^n)$. In [6] Strauss shows that if $\lambda > 0$ and $p > 1$, Problem (1.1)–(1.2) has at least one global weak solution $u(t) \in L^\infty(\mathbf{R}; H^1(\mathbf{R}^n) \cap L^{p+1}(\mathbf{R}^n))$ for any $u_0 \in H^1(\mathbf{R}^n) \cap L^{p+1}(\mathbf{R}^n)$ (see also [5]). In [10] M. Tsutsumi and N. Hayashi discuss the unique global existence of classical solutions for (1.1)–(1.2) with $\lambda > 0$ (see also Pecher and von Wahl [4]). Furthermore, in [9] M. Tsutsumi discusses the unique global solution in $\mathcal{S}(\mathbf{R}^n)$ or in the weighted Sobolev space for (1.1)–(1.2) with $\lambda > 0$. In almost all of previous papers the solution of (1.1)–(1.2) has been constructed in a space not larger than the energy space, that is, $H^1(\mathbf{R}^n)$, because the proofs in almost all of previous papers are based on the energy conservation law. However, in [7] Strauss constructs the wave operators from $L^2(\mathbf{R}^n)$ to $L^2(\mathbf{R}^n)$ for the equation (1.1) with $p=1+4/n$ (see [7, Theorem 5]). His results are almost equivalent to the construction in $L^2(\mathbf{R}^n)$ of unique local solutions for (1.1)–(1.2) with $p=1+4/n$. In this paper we prove that when $1 < p < 1 + 4/n$, we can construct the unique global solution of (1.1)–(1.2) for u_0 in $L^2(\mathbf{R}^n)$ (but possibly not in $H^1(\mathbf{R}^n)$). Such a solution is

called an " L^2 -solution". Furthermore, we show that when $1 < p < 1 + 4/n$, the solution operator of the evolution equation (1.1) constitutes a strongly continuous nonlinear operator group in $L^2(\mathbf{R}^n)$. Our proof is based on the L^2 -norm conservation law and the dispersive effect of solutions (see, e.g., Lemma 2.2).

Our main theorem in this paper is the following.

Theorem 1.1. *Assume that $1 < p < 1 + 4/n$. Then, for any $u_0 \in L^2(\mathbf{R}^n)$ and any $t_0 \in \mathbf{R}$ there exists a unique global solution $u(t)$ of (1.1)–(1.2) such that*

$$(1.3) \quad u(t) \in C(\mathbf{R}; L^2(\mathbf{R}^n)) \cap L'_{\text{loc}}(\mathbf{R}; L^{p+1}(\mathbf{R}^n)),$$

$$(1.4) \quad u(t) = U(t-t_0)u_0 - i \int_{t_0}^t U(t-\tau)f(u(\tau))d\tau, \quad t \in \mathbf{R},$$

$$(1.5) \quad \|u(t)\|_{L^2(\mathbf{R}^n)} = \|u_0\|_{L^2(\mathbf{R}^n)}, \quad t \in \mathbf{R},$$

where $r = \frac{4(p+1)}{n(p-1)}$, $U(t) = e^{i\Delta t}$, $f(z) = \lambda|z|^{p-1}z$ ($z \in \mathbf{C}$) and the integral in (1.4) is the Bochner integral in $H^{-1}(\mathbf{R}^n)$. Furthermore, let u_{0j} , $j=1, 2, \dots$, and u_0 be such that $u_{0j}, u_0 \in L^2(\mathbf{R}^n)$ and $u_{0j} \rightarrow u_0$ in $L^2(\mathbf{R}^n)$ ($j \rightarrow \infty$). Let $u_j(t)$ and $u(t)$ be the solutions of (1.1) with $u_j(t_0) = u_{0j}$ and $u(t_0) = u_0$, respectively. Then, for each $t \in \mathbf{R}$

$$(1.6) \quad u_j(t) \longrightarrow u(t) \quad \text{in } L^2(\mathbf{R}^n) \quad (j \rightarrow \infty).$$

By Theorem 1.1 we can define the solution operator of the evolution equation (1.1) as a mapping from $L^2(\mathbf{R}^n)$ to $L^2(\mathbf{R}^n)$, when $1 < p < 1 + 4/n$. We denote it by $S(t)$. The following result is an immediate consequence of Theorem 1.1.

Corollary 1.2. *Assume that $1 < p < 1 + 4/n$. Then, $\{S(t); -\infty < t < +\infty\}$ is a strongly continuous nonlinear operator group in $L^2(\mathbf{R}^n)$. That is, $S(t)$ is a homeomorphism from $L^2(\mathbf{R}^n)$ to $L^2(\mathbf{R}^n)$ for each $t \in \mathbf{R}$, and*

$$(1.7) \quad S(t+s) = S(t)S(s), \quad t, s \in \mathbf{R},$$

$$(1.8) \quad S(0) = I,$$

$$(1.9) \quad S(h)v \longrightarrow v \quad \text{in } L^2(\mathbf{R}^n) \quad (h \rightarrow 0), \quad v \in L^2(\mathbf{R}^n),$$

where I is the identity operator from $L^2(\mathbf{R}^n)$ to $L^2(\mathbf{R}^n)$.

Our plan in this paper is as follows. In Section 2 we summarize several lemmas needed for the proof of Theorem 1.1. In Section 3 we give a proof of Theorem 1.1.

We conclude this section with several notations given. We abbreviate $L^q(\mathbf{R}^n)$ and $H^m(\mathbf{R}^n)$ to L^q and H^m , respectively. (\cdot, \cdot) denotes the scalar product in L^2 . For a closed interval I in \mathbf{R} and a Hilbert space H we denote the set of all weakly

continuous functions from I to H by $C_w(I; H)$. Let $U(t) = e^{iAt}$ be the evolution operator of the free Schrödinger equation. We put $f(z) = \lambda|z|^{p-1}z$ for $z \in \mathbf{C}$. Let $h(x)$ be an even and positive function in $C_0^\infty(\mathbf{R}^n)$ with $\|h\|_{L^1} = 1$. We put $h_j(x) = j^n h(jx)$ for each positive integer j . $*$ denotes the convolution with respect to spatial variables. In the course of calculations below various constants will be simply denoted by C . In particular, $C = C(*, \dots, *)$ will denote a constant depending only on the quantities appearing in parentheses.

§ 2. Lemmas.

In this section we summarize several results needed for the proof of Theorem 1.1.

For $U(t)$ we have the following two lemmas.

Lemma 2.1. *Let q and r be positive numbers such that $1/q + 1/r = 1$ and $2 \leq q \leq \infty$. For any $t \neq 0$, $U(t)$ is a bounded operator from L^r to L^q satisfying*

$$(2.1) \quad \|U(t)v\|_{L^q} \leq (4\pi|t|)^{n/q-n/2} \|v\|_{L^r}, \quad v \in L^r, \quad t \neq 0,$$

and for any $t \neq 0$, the map $t \rightarrow U(t)$ is strongly continuous. For $q=2$, $U(t)$ is unitary and strongly continuous for all $t \in \mathbf{R}$.

Lemma 2.2. *Let q and r be positive numbers such that $1 \leq q-1 < \alpha(n)$ and $(n/2 - n/q)r = 2$. Then,*

$$(2.2) \quad \|U(\cdot)v\|_{L^r(\mathbf{R}; L^q)} \leq C \|v\|_{L^2},$$

where $C = C(n, q)$.

Lemma 2.1 is well known (see, e.g., [2, Lemma 1.2]). For Lemma 2.2, see Strichartz [8, Corollary 1 in §3] and Ginibre and Velo [3, Proposition 7].

Furthermore, we need the following two lemmas.

Lemma 2.3. *Let I be an open interval in \mathbf{R} . Let $1 < q, r < \infty$ and $a, b > 0$. We put*

$$M = \{v(t) \in L^\infty(I; L^2) \cap L^r(I; L^q); \\ \|v(t)\|_{L^2} \leq a, \quad \text{a.e. } t \in I, \\ \|v\|_{L^r(I; L^q)} \leq b\}.$$

Then M is a closed subset in $L^r(I; L^q)$.

Proof. Let $\{v_j(t)\} \subset M$ and $v(t) \in L^r(I; L^q)$ be such that $\|v_j - v\|_{L^r(I; L^q)} \rightarrow 0$ ($j \rightarrow \infty$). Then, since $\{v_j(t)\}$ is bounded in $L^\infty(I; L^2)$, we can choose a subsequence $\{v_{j_k}(t)\} \subset \{v_j(t)\}$ and $w(t) \in L^\infty(I; L^2)$ such that

$$(2.3) \quad v_{j'}(t) \longrightarrow w(t) \quad \text{*}-\text{weakly in } L^\infty(I; L^2).$$

Since by the definitions of $\{v_{j'}(t)\}$ and $v(t)$ and (2.3) we have

$$(2.4) \quad v_{j'}(t) \longrightarrow v(t) \quad \text{in } \mathcal{D}'(I \times \mathbf{R}^n) \quad (j' \rightarrow \infty),$$

$$(2.5) \quad v_{j'}(t) \longrightarrow w(t) \quad \text{in } \mathcal{D}'(I \times \mathbf{R}^n) \quad (j' \rightarrow \infty),$$

we conclude by the uniqueness of limit that $v(t) = w(t) \in L^\infty(I; L^2) \cap L^r(I; L^q)$. Moreover, we have by (2.3) and the definitions of $\{v_{j'}(t)\}$ and $v(t)$

$$(2.6) \quad \|v\|_{L^\infty(I; L^2)} \leq \liminf_{j' \rightarrow \infty} \|v_{j'}\|_{L^\infty(I; L^2)} \leq a,$$

$$(2.7) \quad \|v\|_{L^r(I; L^q)} = \lim_{j' \rightarrow \infty} \|v_{j'}\|_{L^r(I; L^q)} \leq b,$$

which imply that $v(t) \in M$. This completes the proof of Lemma 2.3. Q. E. D.

Lemma 2.4. *Let T_1 and T_2 be constants with $T_1 < T_2$. Assume that $v(t) \in C([T_1, T_2]; H^{-1})$ and for some $K > 0$*

$$(2.8) \quad \|v(t)\|_{L^2} \leq K, \quad \text{a.e. } t \in [T_1, T_2].$$

Then, $v(t) \in C_w([T_1, T_2]; L^2)$ and (2.8) holds for all $t \in [T_1, T_2]$.

Proof. We first prove that (2.8) holds for all $t \in [T_1, T_2]$. For that purpose, we assume otherwise and derive the contradiction. We assume that (2.8) does not hold for some $t_0 \in [T_1, T_2]$. Then, by (2.8) we can choose $\{t_n\} \subset [T_1, T_2]$ such that $t_n \rightarrow t_0$ ($n \rightarrow \infty$) and (2.8) holds for all t_n . Therefore, we can choose a subsequence $\{t_{n'}\} \subset \{t_n\}$ and $w \in L^2$ such that

$$(2.9) \quad v(t_{n'}) \longrightarrow w \quad \text{weakly in } L^2 \quad (n' \rightarrow \infty).$$

On the other hand, since $v(t) \in C([T_1, T_2]; H^{-1})$, we have

$$(2.10) \quad v(t_{n'}) \longrightarrow v(t_0) \quad \text{in } H^{-1} \quad (n' \rightarrow \infty).$$

(2.9) and (2.10) give us

$$(2.11) \quad v(t_{n'}) \longrightarrow w \quad \text{in } \mathcal{D}'(\mathbf{R}^n) \quad (n' \rightarrow \infty),$$

$$(2.12) \quad v(t_{n'}) \longrightarrow v(t_0) \quad \text{in } \mathcal{D}'(\mathbf{R}^n) \quad (n' \rightarrow \infty).$$

By (2.11–12) and the uniqueness of limit we obtain $v(t_0) = w \in L^2$. Moreover,

$$(2.13) \quad \|v(t_0)\|_{L^2} \leq \liminf_{n' \rightarrow \infty} \|v(t_{n'})\|_{L^2} \leq K,$$

which contradicts the assumption of t_0 . Accordingly, we conclude that (2.8) holds for all $t \in [T_1, T_2]$.

We next prove that $v(t) \in C_w([T_1, T_2]; L^2)$. For any $t_0 \in [T_1, T_2]$, let $\{t_n\}$

be an arbitrary sequence $\subset [T_1, T_2]$ such that $t_n \rightarrow t_0$ ($n \rightarrow \infty$). Since $v(t) \in C([T_1, T_2]; H^{-1})$, we have

$$(2.14) \quad (v(t_n), \psi) \longrightarrow (v(t_0), \psi) \quad (n \rightarrow \infty)$$

for $\psi \in H^1$. By the boundedness in L^2 of $\{v(t_n)\}$ and (2.14) we obtain

$$(2.15) \quad v(t_n) \longrightarrow v(t_0) \quad \text{weakly in } L^2 \quad (n \rightarrow \infty),$$

which implies the weak continuity in L^2 of $v(t)$ at $t = t_0$. Since t_0 is an arbitrary point in $[T_1, T_2]$, we conclude that $v(t) \in C_w([T_1, T_2]; L^2)$. Q. E. D.

Finally we describe the results of Ginibre and Velo [2] concerning the unique global existence of weak solutions for (1.1)–(1.2). We first formulate this problem precisely. We consider the integral equation

$$(2.16) \quad u(t) = U(t-t_0)u_0 - i \int_{t_0}^t U(t-\tau)f(u(\tau))d\tau,$$

as the integral version of the initial value problem (1.1)–(1.2). For (2.16) we have the following proposition (see, e.g., [2, Theorem 3.1]).

Proposition 2.5. *Assume that $1 < p < 1 + 4/n$. Then, for any $t_0 \in \mathbf{R}$ and any $u_0 \in H^1$ there exists a unique solution $u(t)$ of (2.16) such that*

$$(2.17) \quad u(t) \in C(\mathbf{R}; H^1),$$

$$(2.18) \quad \|u(t)\|_{L^2} = \|u_0\|_{L^2}, \quad t \in \mathbf{R}.$$

§3. Proof of Theorem 1.1.

In this section we give a proof of Theorem 1.1. By I_t and \bar{I}_t we denote an open interval $(t_0 - t, t_0 + t)$ and a closed interval $[t_0 - t, t_0 + t]$, respectively, for $t \geq 0$. Let $r = \frac{4(p+1)}{n(p-1)}$ throughout this section.

We first prove the following result concerning the unique local existence of L^2 -solutions for (1.1)–(1.2).

Lemma 3.1. *Assume that $1 < p < 1 + 4/n$. Then, for any $t_0 \in \mathbf{R}$ and any $u_0 \in L^2$ there exists a $T = T(p, n, \lambda, \|u_0\|_{L^2}) > 0$ such that Problem (1.1)–(1.2) has a unique local solution $u(t)$:*

$$(3.1) \quad u(t) \in C(\bar{I}_T; L^2) \cap L^r(I_T; L^{p+1}),$$

$$(3.2) \quad u(t) = U(t-t_0)u_0 - i \int_{t_0}^t U(t-\tau)f(u(\tau))d\tau, \quad t \in \bar{I}_T,$$

where the integral in (3.2) is the Bochner integral in H^{-1} . Furthermore, the solution $u(t)$ satisfies

$$(3.3) \quad \|u(t)\|_{L^2} = \|u_0\|_{L^2}, \quad t \in \bar{I}_T.$$

Proof. We consider the following integral equation:

$$(3.4) \quad u_j(t) = U(t-t_0)h_j * u_0 - i \int_{t_0}^t U(t-\tau)f(u_j(\tau))d\tau, \quad j = 1, 2, \dots.$$

By Proposition 2.5 we have the unique global solution $u_j(t)$ in $C(\mathbf{R}; H^1)$ of (3.4) for each j . Let $\rho = \|u_0\|_{L^2}$. By δ we denote the constant appearing in (2.2) with $q = p+1$ and $r = \frac{4(p+1)}{n(p-1)}$. We note that δ depends only on n and p . We put

$$(3.5) \quad M = \{v(t) \in L^\infty(I_T; L^2) \cap L^r(I_T; L^{p+1}); \\ \|v\|_{L^\infty(I_T; L^2)} \leq \rho, \|v\|_{L^r(I_T; L^{p+1})} \leq 2\delta\rho\},$$

where T is a small positive constant to be determined later. We note that by Lemma 2.3 M is closed in $L^r(I_T; L^{p+1})$.

We first show that if T is sufficiently small, then

$$(3.6) \quad u_j(t) \in M \quad \text{for all } j.$$

By (2.18) we have

$$(3.7) \quad \|u_j(t)\|_{L^2} = \|h_j * u_0\|_{L^2} \leq \|u_0\|_{L^2}, \quad t \in \bar{I}_T,$$

for all j . We define $\tilde{u}_j^s(t)$ by

$$(3.8) \quad \tilde{u}_j^s(t) = \begin{cases} u_j(t), & t \in I_s, \\ 0, & t \notin I_s, \end{cases}$$

for $s \geq 0$. By (3.4) and (2.1-2) we have for s with $0 \leq s \leq T$

$$(3.9) \quad \|u\|_{L^r(I_s; L^{p+1})} \leq \delta\rho \\ + C \left\| \int_{t_0}^t |t-\tau|^{-n/2+n/(p+1)} \|f(u_j(\tau))\|_{L^{1+1/p}} d\tau \right\|_{L^r(I_s)} \\ \leq \delta\rho + C \left\| \int_{-\infty}^{+\infty} |t-\tau|^{-n/2+n/(p+1)} \|\tilde{u}_j^s(\tau)\|_{L^{p+1}}^p d\tau \right\|_{L^r(\mathbf{R})} \\ \leq \delta\rho + C \|\tilde{u}_j^s\|_{L^{q_1}(\mathbf{R}; L^{p+1})}^p \\ \leq \delta\rho + C \|u_j\|_{L^{q_1}(I_s; L^{p+1})}^p, \quad j = 1, 2, \dots,$$

where $q_1 = \frac{4p(p+1)}{n+4-(n-4)p}$. Here we have used the generalized Young inequality

(see [12, §4 in Chapter IX]) at the third inequality. We note that $1 < q_1 < r$ for $1 < p < 1 + 4/n$. On the other hand, we have by Hölder's inequality

$$(3.10) \quad \|u_j\|_{L^{q_1}(I_s; L^{p+1})} \leq \left(\int_{t_0-s}^{t_0+s} d\tau \right)^{1/q_2} \|u_j\|_{L^r(I_s; L^{p+1})} \\ \leq CT^{1/q_2} \|u_j\|_{L^r(I_s; L^{p+1})}, \quad 0 \leq s \leq T, \quad j = 1, 2, \dots,$$

where $q_2 = \frac{4p}{n+4-np}$ and $C = C(n, p)$. (3.9) and (3.10) give us

$$(3.11) \quad \|u_j\|_{L^r(I_s; L^{p+1})} \leq \delta\rho + C_0 T^{p/q_2} \|u_j\|_{L^r(I_s; L^{p+1})}^p, \\ 0 \leq s \leq T, \quad j = 1, 2, \dots,$$

where $C_0 = C_0(n, p, \lambda)$. Now we choose $T > 0$ so small that there exists a positive number y satisfying $C_0 T^{p/q_2} y^p + \delta\rho - y < 0$ and $0 < y \leq 2\delta\rho$. For that purpose, it is sufficient to choose $T > 0$ so that

$$(3.12) \quad T < (2C_0(2\delta\rho)^{p-1})^{-q_2/p}.$$

Then we put

$$(3.13) \quad y_0 = \min \{2\delta\rho \geq y > 0; C_0 T^{p/q_2} y^p + \delta\rho - y = 0\}.$$

Putting $X_j(s) = \|u_j\|_{L^r(I_s; L^{p+1})}$, we have by (3.11)

$$(3.14) \quad X_j(s) \leq \delta\rho + C_0 T^{p/q_2} X_j(s)^p, \quad 0 \leq s \leq T,$$

$$(3.15) \quad X_j(0) = 0,$$

for all j . If T is chosen so small that (3.12) holds, then by (3.13) and (3.14–15) we obtain

$$(3.16) \quad X_j(s) \leq y_0 \leq 2\delta\rho, \quad 0 \leq s \leq T, \quad j = 1, 2, \dots.$$

From (3.16) and Fatou's lemma it follows that

$$(3.17) \quad \|u_j\|_{L^r(I_T; L^{p+1})} \leq 2\delta\rho, \quad j = 1, 2, \dots.$$

By (3.7) and (3.17) we obtain (3.6), if T is chosen so small that (3.12) holds. From now on we assume that T is chosen so small that (3.12) holds.

For any j and k , in the same way as (3.9–11) we have by (3.4), Lemmas 2.1, 2.2 and the generalized Young inequality

$$(3.18) \quad \|u_j - u_k\|_{L^r(I_T; L^{p+1})} \leq K(j, k) \\ + C \int_{t_0}^t |t - \tau|^{-n/2+n/(p+1)} \|f(u_j(\tau)) - f(u_k(\tau))\|_{L^{1+1/p}} d\tau \|L^r(I_T)$$

$$\begin{aligned}
&\leq K(j, k) + C \left\| \int_{t_0}^t |t-\tau|^{-n/2+n/(p+1)} \right. \\
&\quad \times (\|u_j(\tau)\|_{L^{p+1}}^{p-1} + \|u_k(\tau)\|_{L^{p+1}}^{p-1}) \|u_j(\tau) - u_k(\tau)\|_{L^{p+1}} d\tau \left. \right\|_{L^r(I_T)} \\
&\leq K(j, k) + C (\|u_j\|_{L^{q_1}(I_T; L^{p+1})}^{p-1} + \|u_k\|_{L^{q_1}(I_T; L^{p+1})}^{p-1}) \\
&\quad \times \|u_j - u_k\|_{L^{q_1}(I_T; L^{p+1})} \\
&\leq K(j, k) + \bar{C}_0 T^{p/q_2} (\|u_j\|_{L^r(I_T; L^{p+1})}^{p-1} + \|u_k\|_{L^r(I_T; L^{p+1})}^{p-1}) \\
&\quad \times \|u_j - u_k\|_{L^r(I_T; L^{p+1})} \\
&\leq K(j, k) + \bar{C}_0 T^{p/q_2} \cdot 2(2\delta\rho)^{p-1} \|u_j - u_k\|_{L^r(I_T; L^{p+1})},
\end{aligned}$$

where $K(j, k) = \delta \|h_j * u_0 - h_k * u_0\|_{L^2}$, $q_1 = \frac{4p(p+1)}{n+4-(n-4)p}$, $q_2 = \frac{4p}{n+4-np}$ and $\bar{C}_0 = \bar{C}_0(n, p, \lambda)$. We have used the generalized Young inequality at the third inequality and have used Hölder's inequality with $1/q_2 + q_1/r = 1$ at the last inequality but one. If we choose T so small in (3.18) that

$$(3.19) \quad \bar{C}_0 T^{p/q_2} \cdot 2(2\delta\rho)^{p-1} \leq \frac{1}{2},$$

then we have by (3.18)

$$(3.20) \quad \|u_j - u_k\|_{L^r(I_T; L^{p+1})} \leq 2K(j, k)$$

for all j and k . Since $K(j, k) \rightarrow 0$ ($j, k \rightarrow \infty$), we have by (3.20)

$$(3.21) \quad \|u_j - u_k\|_{L^r(I_T; L^{p+1})} \longrightarrow 0 \quad (j, k \rightarrow \infty),$$

if T is chosen so small that (3.19) holds. Furthermore, we have by (3.4), (3.21), the Sobolev imbedding theorem and Hölder's inequality

$$\begin{aligned}
(3.22) \quad & |(u_j(t) - u_k(t), \psi)| \\
& \leq \|h_j * u_0 - h_k * u_0\|_{L^2} \|\psi\|_{L^2} \\
& \quad + \left| \int_{t_0}^t (U(t-\tau) \{f(u_j(\tau)) - f(u_k(\tau))\}, \psi) d\tau \right| \\
& \leq \|h_j * u_0 - h_k * u_0\|_{L^2} \|\psi\|_{L^2} \\
& \quad + \int_{t_0-T}^{t_0+T} \|f(u_j(\tau)) - f(u_k(\tau))\|_{L^{1+1/p}} \|U(\tau-t)\psi\|_{L^{p+1}} d\tau \\
& \leq \|h_j * u_0 - h_k * u_0\|_{L^2} \|\psi\|_{L^2} \\
& \quad + C \|\psi\|_{H^1} \int_{t_0-T}^{t_0+T} (\|u_j(\tau)\|_{L^{p+1}}^{p-1} + \|u_k(\tau)\|_{L^{p+1}}^{p-1}) \\
& \quad \times \|u_j(\tau) - u_k(\tau)\|_{L^{p+1}} d\tau
\end{aligned}$$

$$\begin{aligned} &\leq \|h_j * u_0 - h_k * u_0\|_{L^2} \|\psi\|_{L^2} \\ &\quad + CT^{q_3} \|\psi\|_{H^1} \cdot 2(2\delta\rho)^{p-1} \|u_j - u_k\|_{L^r(I_T; L^{p+1})} \\ &\quad \longrightarrow 0 \quad (j, k \rightarrow \infty) \quad \text{uniformly in } t \in \bar{I}_T, \end{aligned}$$

for $\psi \in H^1$, where $q_3 = \frac{4 + (n+4)p - np^2}{4(p+1)}$. We note that $q_3 > 0$ for $1 < p < 1 + 4/n$. Since by Proposition 2.5 $u_j(t) \in C(\bar{I}_T; H^{-1})$ for any j , (3.22) implies that $\{u_j(t)\}$ is the Cauchy sequence in $C(\bar{I}_T; H^{-1})$.

Therefore, by (3.6), (3.21), (3.22) and Lemma 2.3 we obtain the solution $u(t)$ of (1.1)–(1.2) such that

$$(3.23) \quad u(t) \in L^\infty(I_T; L^2) \cap L^r(I_T; L^{p+1}) \cap C(\bar{I}_T; H^{-1}),$$

$$(3.24) \quad u(t) = U(t-t_0)u_0 - i \int_{t_0}^t U(t-\tau)f(u(\tau))d\tau, \quad t \in \bar{I}_T,$$

$$(3.25) \quad \|u(t)\|_{L^2} \leq \|u_0\|_{L^2}, \quad \text{a.e. } t \in I_T$$

$$(3.26) \quad u_j(t) \longrightarrow u(t) \quad \text{in } L^r(I_T; L^{p+1}) \quad \text{and in } C(\bar{I}_T; H^{-1}) \quad (j \rightarrow \infty),$$

where T is a positive constant with (3.12) and (3.19) and the integral in (3.24) is the Bochner integral in H^{-1} . (3.23), (3.25) and Lemma 2.4 imply that

$$(3.27) \quad u(t) \in C_w(\bar{I}_T; L^2)$$

and that for all $t \in \bar{I}_T$ (3.25) holds.

We next show that the solution satisfying (3.23) and (3.24) is unique. Let $u(t)$ and $v(t)$ be two solutions satisfying (3.23) and (3.24) with the same initial datum. We put $t_1 = \sup\{t \in [0, T]; u(s) = v(s) \text{ on } \bar{I}_t\}$. If $t_1 = T$, then $u(t) = v(t)$ on \bar{I}_T , which is the desired result. If $t_1 < T$, in the same way as (3.18) we have by (3.24) and the assumption of t_1

$$\begin{aligned} (3.28) \quad &\|u - v\|_{L^r(I_{t_1, t_2}; L^{p+1})} = \|u - v\|_{L^r(I_{t_2}; L^{p+1})} \\ &\leq C(t_1 - t_2)^{p/q_2} (\|u\|_{L^r(I_{t_1, t_2}; L^{p+1})}^{p-1} + \|v\|_{L^r(I_{t_1, t_2}; L^{p+1})}^{p-1}) \\ &\quad \times \|u - v\|_{L^r(I_{t_1, t_2}; L^{p+1})}, \quad t_2 \in (t_1, T], \end{aligned}$$

where $q_2 = \frac{4p}{n+4-np}$, $C = C(n, q, p)$ and $I_{t_1, t_2} = (-t_2, -t_1) \cup (t_1, t_2)$. We can choose t_2 such that $t_2 > t_1$ and the coefficient of $\|u - v\|_{L^r(I_{t_1, t_2}; L^{p+1})}$ is smaller than $1/2$ in the right hand side of (3.28). Then,

$$(3.29) \quad \|u - v\|_{L^r(I_{t_1, t_2}; L^{p+1})} \leq 0,$$

which implies that $u(t) = v(t)$ on \bar{I}_{t_1, t_2} . This contradicts the assumption of t_1 .

Therefore, $u(t) = v(t)$ for $t \in \bar{I}_T$.

Thus, for any $t_1 \in \bar{I}_T$ we can uniquely solve (1.1)–(1.2) in the time interval $[t_1 - T, t_1 + T]$ with the initial time t_0 and the initial datum u_0 replaced by t_1 and $u(t_1)$, respectively, where T is the same as in the case of the initial time t_0 and the initial datum u_0 . Therefore, reversing the roles of 0 and t , we obtain the reverse inequality to (3.25) for all $t \in \bar{I}_T$. Accordingly, we obtain (3.3). (3.3) and (3.27) give us

$$(3.30) \quad u(t) \in C(\bar{I}_T; L^2).$$

This completes the proof of Lemma 3.1.

Q. E. D.

We are now in a position to prove Theorem 1.1.

Proof of Theorem 1.1. The unique global existence of L^2 -solutions for (1.1)–(1.2) follows directly from Lemma 3.1, which shows the unique local solvability in L^2 of (1.1)–(1.2) and the a priori bound of the L^2 -norm of L^2 -solutions.

It remains only to prove the continuous dependence of L^2 -solutions on the initial data. Let u_{0j} , $j = 1, 2, \dots$, and u_0 be such that $u_{0j}, u_0 \in L^2$ and $u_{0j} \rightarrow u_0$ in L^2 ($j \rightarrow \infty$). Let $u_j(t)$ and $u(t)$ be the global L^2 -solutions of (1.1) with $u_j(t_0) = u_{0j}$ and $u(t_0) = u_0$, respectively. We put $\rho = \sup \{ \|u_0\|_{L^2}, \|u_{0j}\|_{L^2}, j = 1, 2, \dots \}$. For this ρ , let $T > 0$ be defined as in (3.12) and (3.19). Then, by using the same argument as in the proof of Lemma 3.1 we have

$$(3.31) \quad u_j(t) \longrightarrow u(t) \quad \text{in } C(\bar{I}_t; H^{-1}) \quad (j \rightarrow \infty)$$

(see, e.g., (3.22) and (3.26)). Particularly, we have

$$(3.32) \quad (u_j(t), \psi) \longrightarrow (u(t), \psi), \quad t \in \bar{I}_t$$

for $\psi \in H^1$. By (3.32) and the boundedness in L^2 of $\{u_j(t)\}$ we obtain

$$(3.33) \quad u_j(t) \longrightarrow u(t) \quad \text{weakly in } L^2 \quad (j \rightarrow \infty), \quad t \in \bar{I}_T.$$

On the other hand, we have by (1.5)

$$(3.34) \quad \|u_j(t)\|_{L^2} \longrightarrow \|u(t)\|_{L^2} \quad (j \rightarrow \infty), \quad t \in \mathbf{R}.$$

Therefore, (3.33) and (3.34) give us

$$(3.35) \quad u_j(t) \longrightarrow u(t) \quad \text{in } L^2 \quad (j \rightarrow \infty)$$

for each $t \in \bar{I}_T$. The length of T is determined only by n , p , λ and ρ (see (3.12) and (3.19)). By the L^2 -norm conservation law (see (1.5) and (3.3)) we see that $\sup \{ \|u(t)\|_{L^2}, \|u_j(t)\|_{L^2}, j = 1, 2, \dots \}$ is constant for $t \in \mathbf{R}$. Accordingly, we use the above argument with the initial time t_0 and the initial data $u_0, u_{0j}, j = 1, 2, \dots$,

replaced by $t_0 + T$ and $u(t_0 + T)$, $u_j(t_0 + T)$, $j = 1, 2, \dots$, or by $t_0 - T$ and $u(t_0 - T)$, $u_j(t_0 - T)$, $j = 1, 2, \dots$, respectively, to obtain (3.35) for each $t \in \bar{I}_{2T}$. Repeating this procedure, we obtain (1.6) for each $t \in \mathbf{R}$. This completes the proof of Theorem 1.1. Q. E. D.

Acknowledgement. The author would like to express his deep gratitude to Professor S. T. Kuroda for his helpful conversation and constant encouragement.

References

- [1] Baillon, J. B., Cazenave, T. and Figueira, M., Équation de Schrödinger nonlinéaire, C. R. Acad. Sci., Paris, **284** (1977), 867–872.
- [2] Ginibre, J. and Velo, G., On a class of nonlinear Schrödinger equations. I: The Cauchy problem, J. Funct. Anal., **32** (1979), 1–32.
- [3] ——— and ———, Theorie de la diffusion dans l'espace d'énergie pour une classe d'équations de Schrödinger non lineaire, C. R. Acad. Sci., Paris, **298** (1984), 137–140.
- [4] Pecher, H. and von Wahl, W., Time dependent nonlinear Schrödinger equations, Manuscripta Math., **27** (1979), 125–157.
- [5] Strauss, W. A., On weak solutions of semi-linear hyperbolic equations, An. Acad. Brasil. Cienc., **42** (1970), 645–651.
- [6] Strauss, W. A., The nonlinear Schrödinger equation, in "Contemporary Developments in Continuum Mechanics and Partial Differential Equations", North-Holland, Amsterdam-New York-Oxford, 1978.
- [7] Strauss, W. A., Everywhere defined wave operators, in "Nonlinear Evolution Equations", pp. 85–102, Academic Press, New York, 1978.
- [8] Strichartz, R. S., Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations, Duke Math. J., **44** (1977), 705–714.
- [9] Tsutsumi, M., Weighted Sobolev spaces and rapidly decreasing solutions of some nonlinear dispersive wave equations, J. Differential Equations, **42** (1981), 260–281.
- [10] Tsutsumi, M. and Hayashi, N., Classical solutions of nonlinear Schrödinger equations in higher dimensions, Math. Z., **177** (1981), 217–234.
- [11] Tsutsumi, Y., Global existence and asymptotic behavior of solutions for nonlinear Schrödinger equations, Doctor thesis, University of Tokyo, 1985.
- [12] Reed, M. and Simon, B., *Methods of Modern Mathematical Physics, Vol. II: Fourier Analysis and Self-adjointness*, Academic Press, New York, 1975.

nuna adreso:
Faculty of Integrated Arts and Sciences
Hiroshima University
Higashisenda-machi, Naka-ku
Hiroshima 730, Japan

(Ricevita la 8-an de julio, 1985)