

## Cauchy Problem for Some Degenerate Abstract Differential Equations of Sobolev Type

By

Tomomi KOJO and Masayoshi TSUTSUMI

(Shibaura Institute of Technology and Waseda University, Japan)

### 1. Introduction

Let  $H$  be a Hilbert space over the reals. We shall consider an ordinary differential equation in  $H$ :

$$(E) \quad B \frac{du}{dt}(t) + Au(t) \ni f(t), \quad t \geq 0,$$

together with the initial condition

$$(IC) \quad B^{1/2}u(0) = B^{1/2}u_0.$$

Here  $A$  is a nonlinear  $m$ -accretive operator in  $H$  and  $B$  is a nonnegative selfadjoint operator in  $H$ .

Equations of the form (E) appear in various physical problems including the propagation of long waves of small amplitude [3], the heat conduction involving two temperatures [7] and soil mechanics [2] and are called pseudo-parabolic equations (since they are parabolic equations when  $B = I$ ) or equations of Sobolev type [12].

In the previous works, the authors have established the existence-uniqueness theorem of strong solutions of the initial value problem for (E) under various conditions on  $A$  and  $B$ :

- i)  $D(B) \subset D(A)$  and  $B$  has the bounded inverse [13],
- ii)  $D(B) \subset D(A)$  and  $B$  is not necessarily invertible [14],
- iii)  $D(A) \subset D(B)$  and  $B$  has the bounded inverse [8].

The purpose of this paper is to construct a result for (E), (IC) under the case

- iv)  $B$  is not necessarily invertible (and no inclusion relation between  $D(A)$  and  $D(B)$  is assumed),

and apply it to the initial-boundary value problem for some nonlinear partial differential equations.

The pseudo-parabolic equation is investigated, in the abstract frame work, by many authors under various conditions on  $A$  and  $B$  which are different from ours [5], [6], [9], [11], [15] (and the references therein).

## 2. Notation and a result

Let  $H$  be a real Hilbert space, the inner product and the norm in  $H$  denoted by  $(\cdot, \cdot)$  and  $\|\cdot\|$  respectively.

Let  $A$  be a nonlinear multivalued operator from  $H$  into itself. An operator  $A: H \rightarrow H$  with domain  $D(A) = \{u; Au \neq \emptyset\}$  and range  $R(A) = \cup\{Au; u \in D(A)\}$  is said to be *accretive* if

$$(v_1 - v_2, u_1 - u_2) \geq 0 \quad \text{for every } v_j \in Au_j \quad (u_j \in D(A), j = 1, 2).$$

An accretive operator  $A$  is said to be *m-accretive* if  $R(I + A) = H$ . For each integer  $n > 0$  the *Yosida approximation*  $A_n$  of an *m-accretive* operator  $A$  is defined by

$$A_n = n(I - (I + n^{-1}A)^{-1}).$$

It is well known that  $A_n$  is *m-accretive* on  $H$  and Lipschitz continuous with  $n$  as Lipschitz constant. For brevity of notation, we shall denote  $(I + n^{-1}A)^{-1}$  by  $J_n^A$ . From the definition

$$(\dagger) \quad A_n u \in A J_n^A u \quad \text{for every } u \in H.$$

We shall denote by  $\Phi$  the set of all lower semicontinuous (l.s.c.) convex functions from  $H$  into  $(-\infty, +\infty]$ , not identically  $+\infty$ . For  $\varphi \in \Phi$ , let  $D(\varphi) = \{u \in H; \varphi(u) < +\infty\}$  and denote by  $\partial\varphi$  the *subdifferential* of  $\varphi$ :

$$\partial\varphi(u) = \{\zeta \in H; \varphi(u) - \varphi(v) \leq (\zeta, u - v) \text{ for every } v \in H\}$$

with  $D(\partial\varphi) = \{u \in H; \partial\varphi(u) \neq \emptyset\}$ . It is well known that  $\partial\varphi$  is *m-accretive* in  $H$  and  $D(\partial\varphi)$  is dense in  $D(\varphi)$ . We refer to [1] and [4] for the properties of *m-accretive* operators in a Hilbert space.

By  $AC([0, T]; H)$  we denote the space of all  $H$ -valued strongly absolutely continuous functions on  $[0, T]$ . For other function spaces, we shall employ the usual notation [10].

**Definition 1.** An  $H$ -valued function  $u(t)$  is called a *strong solution* of (E), (IC) if

- 1)  $u \in AC([\delta, T]; H)$  ( $\forall \delta > 0$ ),  $\frac{du}{dt}(t) \in D(B)$  a.e.  $(0, T)$ ,
- 2)  $u(t) \in D(A)$  a.e.  $(0, T)$  and there exists a  $\xi(t) \in Au(t)$  such that

$$B \frac{du}{dt}(t) + \xi(t) = f(t) \quad \text{a.e. } (0, T),$$

- 3)  $B^{1/2}u \in AC([0, T]; H)$  and  $B^{1/2}u(t)$  satisfies (IC).

*Remark 1.* If in addition  $Bu \in AC([0, T]; H)$  then

$$\frac{d}{dt}(Bu)(t) = B \frac{du}{dt}(t) \quad \text{a.e. } (0, T).$$

To establish the existence-uniqueness theorem for (E), (IC), we shall assume the followings.

(A.1)  $A = \partial\varphi$ , where  $\varphi \in \Phi$  and for every  $u \in D(\varphi)$

$$\varphi(u) \geq a\|u\|^2 - a' \quad (a, a' > 0).$$

(A.2) For every  $\xi \in Au$  and  $\eta \in Av$  ( $u, v \in D(A)$ ) there exists a constant  $d > 0$  such that

$$(\xi - \eta, u - v) \geq d\|u - v\|^2.$$

(A.3)  $B$  is a nonnegative selfadjoint operator in  $H$ .

(A.4) For every  $u, v \in D(B)$  and integer  $n > 0$

$$(A_n u - A_n v, B(u - v)) \geq 0.$$

*Remark 2.* Note that

$$(\xi - \eta, B(u - v)) \geq 0 \quad \text{for every } \xi \in Au \text{ and } \eta \in Av \text{ (} u, v \in D(A) \cap D(B)\text{)}$$

implies (A.4) because

$$\begin{aligned} (A_n u - A_n v, B(u - v)) &= \frac{1}{n} \|B^{1/2}(A_n u - A_n v)\|^2 \\ &\quad + (A_n u - A_n v, B(J_n^A u - J_n^A v)) \geq 0. \end{aligned}$$

Here we have used the selfadjointness of  $B$ , the identity

$$(\#) \quad v = n^{-1} A_n v + J_n^A v \quad (v \in H)$$

and  $(\dagger)$ .

**Theorem 1.** Assume that (A.1) ~ (A.4) are satisfied. Then for every  $0 < T < +\infty$ ,  $f \in W^{1,2}(0, T; H)$  and  $u_0 \in D(A) \cap D(B)$  there exists one and only one strong solution  $u(t)$  of (E), (IC) such that

$$u, \xi(\in Au) \in L^2(0, T; H),$$

$$\sqrt{t}(du/dt) \in L^2(0, T; H), \quad (t > 0)$$

and

$$B^{1/2}u, Bu \in AC([0, T]; H), \quad (d/dt)(B^{1/2}u), (d/dt)(Bu) \in L^2(0, T; H).$$

### 3. Proof of Theorem 1

1°. *Uniqueness.* From (A.2) and (A.3) uniqueness of solutions can be obtained by the standard procedure.

2°. *Existence.* Consider an approximate equation for (E):

$$(E_n) \quad \left(\frac{1}{n} + B\right) u_n'(t) + A_n u_n(t) = f(t), \quad t \geq 0,$$

together with

$$(IC_n) \quad u_n(0) = u_0.$$

Here  $' = d/dt$  and for each integer  $n > 0$ ,  $A_n$  is the Yosida approximation of  $A$ . Since the mapping  $v \mapsto (n^{-1} + B)^{-1} A_n v$  from  $H$  into  $H$  is Lipschitz continuous, the theory of ordinary differential equations in Hilbert space yields that for every  $0 < T < +\infty$ ,  $f \in L^2(0, T; H)$  and integer  $n > 0$  there exists a unique function  $u_n(\cdot) \in C^1([0, T]; D(B))$  which satisfies  $(E_n)$  and  $(IC_n)$ . We shall prove that  $u_n$  converges to a solution  $u$  of (E), (IC) as  $n$  tends to infinity. To this end some a priori estimates are necessary.

i) Taking the inner product of the both sides of  $(E_n)$  by  $u_n(t)$  we have

$$(3.1) \quad \frac{1}{2n} \frac{d}{dt} \|u_n(t)\|^2 + \frac{1}{2} \frac{d}{dt} \|B^{1/2} u_n(t)\|^2 + (A_n u_n(t), u_n(t)) = (f(t), u_n(t)).$$

For simplicity, for a function  $v(t) \in H$  we shall suppress a letter  $t$  and denote by  $v$  in several places below. Using the identity (#), (A.2) and (†) we can estimate the terms in (3.1) as

$$(A_n u_n, u_n) = \frac{1}{n} \|A_n u_n\|^2 + (A_n u_n, J_n^A u_n) \geq \frac{1}{n} \|A_n u_n\|^2 + d \|J_n^A u_n\|^2,$$

and

$$\begin{aligned} (f, u_n) &\leq \|f\| \left\| \frac{1}{n} A_n u_n + J_n^A u_n \right\| \\ &\leq \left( \frac{1}{2n} + c \right) \|f\|^2 + \frac{1}{2n} \|A_n u_n\|^2 + \frac{d}{2} \|J_n^A u_n\|^2. \end{aligned}$$

Then we have

$$\begin{aligned} &\frac{1}{2n} \frac{d}{dt} \|u_n(t)\|^2 + \frac{1}{2} \frac{d}{dt} \|B^{1/2} u_n(t)\|^2 + \frac{1}{2n} \|A_n u_n(t)\|^2 + \frac{d}{2} \|J_n^A u_n(t)\|^2 \\ &\leq \left( \frac{1}{2n} + c \right) \|f(t)\|^2. \end{aligned}$$

Here and in the sequel of this paper, by  $c$  we denote various positive constants independent of  $n$  in a certain interval  $[N, +\infty)$  ( $N > 0$ ). Integrating both sides of this inequality over  $(0, t)$  and taking the assumptions on  $u_n(0)$  and  $f(t)$  into

account we see

$$(3.2) \quad \frac{1}{\sqrt{n}} \|u_n(t)\| \leq c \quad \text{for every } t \in [0, T],$$

$$(3.3) \quad \|B^{1/2}u_n(t)\| \leq c \quad \text{for every } t \in [0, T],$$

$$(3.4) \quad \frac{1}{\sqrt{n}} |A_n u_n|_T \leq c$$

and

$$(3.5) \quad |J_n^A u_n|_T \leq c.$$

Here  $|\cdot|_T$  is the norm in  $L^2(0, T; H)$ .

ii) Since  $u_n$  belongs to  $C^1([0, T]; D(B))$  for each integer  $n > 0$

$$(3.6) \quad \left(\frac{1}{n} + B\right)u_n'(0) + A_n u_n(0) = f(0)$$

holds. Taking the inner product of the both sides of (3.6) by  $n^{-1}u_n'(0)$  we have

$$\left\| \frac{1}{n} u_n'(0) \right\|^2 + \left\| \frac{1}{\sqrt{n}} B^{1/2} u_n'(0) \right\|^2 \leq \frac{1}{2} \|f(0) - A_n u_n(0)\|^2 + \frac{1}{2} \left\| \frac{1}{n} u_n'(0) \right\|^2$$

which implies

$$(3.7) \quad \left\| \frac{1}{n} u_n'(0) \right\| \leq c$$

and

$$(3.8) \quad \left\| \frac{1}{\sqrt{n}} B^{1/2} u_n'(0) \right\| \leq c.$$

iii) For every  $t \in [0, T]$ ,  $h > 0$  and integer  $n > 0$

$$(3.9) \quad \left(\frac{1}{n} + B\right)(u_n'(t+h) - u_n'(t)) + A_n u_n(t+h) - A_n u_n(t) = f(t+h) - f(t)$$

holds. Taking the inner product of the both sides of (3.9) by  $B(u_n(t+h) - u_n(t))$  and using (A.3) and (A.4) we have

$$(3.10) \quad \frac{1}{2n} \frac{d}{dt} \|B^{1/2}(u_n(t+h) - u_n(t))\|^2 + \frac{1}{2} \frac{d}{dt} \|B(u_n(t+h) - u_n(t))\|^2 \\ \leq (f(t+h) - f(t), B(u_n(t+h) - u_n(t))).$$

Dividing the both sides of (3.10) by  $h^2$ , integrating it over  $(0, t)$  and letting  $h$

tends to zero we have

$$\begin{aligned} \frac{1}{n} \|B^{1/2}u'_n(t)\|^2 + \|Bu'_n(t)\|^2 &\leq c \int_0^t \|f'(t)\|^2 dt + c \int_0^t \|Bu'_n(t)\|^2 dt \\ &\quad + \|Bu'_n(0)\|^2 + \frac{1}{n} \|B^{1/2}u'_n(0)\|^2 \end{aligned}$$

which implies

$$(3.11) \quad \|Bu'_n(t)\| \leq c \quad \text{for every } t \in [0, T]$$

and

$$(3.12) \quad \frac{1}{\sqrt{n}} \|B^{1/2}u'_n(t)\| \leq c \quad \text{for every } t \in [0, T].$$

Here we have used (3.7), (3.8) and the identity  $Bu'_n(0) = -n^{-1}u'_n(0) - A_nu_n(0) + f(0)$ . From (3.11) and the assumption  $u_0 \in D(B)$  we also have

$$(3.13) \quad \|Bu_n(t)\| \leq c \quad \text{for every } t \in [0, T].$$

iv) Taking the inner product of the both sides of (E<sub>n</sub>) by  $u'_n(t)$  and integrating it over  $(0, t)$  we have

$$\int_0^t \left\| \frac{1}{\sqrt{n}} u'_n(t) \right\|^2 dt + \int_0^t \|B^{1/2}u'_n(t)\|^2 dt + \varphi_n(u_n(t)) = \varphi_n(u_0) + \int_0^t (f(t), u'_n(t)) dt.$$

Here  $\varphi_n$  is defined by  $\varphi_n(v) = (2n)^{-1} \|A_nv\|^2 + \varphi(J_n^A v)$  ( $v \in H$ ) and we have used the fact  $(A_nv(t), v'(t)) = (d/dt)\varphi_n(v(t))$  ([4], Lemme 3.3). In order to get the boundedness of  $\|(1/\sqrt{n})u'_n\|_T$  we can estimate the right hand side of this equality as

$$\begin{aligned} \int_0^t (f(t), u'_n(t)) dt &= (f(t), u_n(t)) - (f(0), u_0) - \int_0^t (f'(t), u_n(t)) dt \\ &\leq \|f(t)\| \left\| \frac{1}{n} A_nu_n(t) + J_n^A u_n(t) \right\| + \|f(0)\| \|u_0\| \\ &\quad + \int_0^t \|f'(t)\| \left\| \frac{1}{n} A_nu_n(t) + J_n^A u_n(t) \right\| dt. \end{aligned}$$

According to (A.1) the first term of the right hand side of this inequality can be estimated from above by

$$\begin{aligned} \|f(t)\| \left\| \frac{1}{n} A_nu_n + J_n^A u_n \right\| &\leq \frac{1}{4n} \|A_nu_n\|^2 + \frac{c}{n} \|f(t)\|^2 + \frac{a}{2} \|J_n^A u_n\|^2 + c \|f(t)\|^2 \\ &\leq \frac{1}{2} \left( \frac{1}{2n} \|A_nu_n\|^2 + \varphi(J_n^A u_n) \right) + \left( \frac{c}{n} + c \right) \|f(t)\|^2 + \frac{a'}{2} \\ &= \frac{1}{2} \varphi_n(u_n(t)) + c \|f(t)\|^2 + c. \end{aligned}$$

Combining them we get by (3.4) and (3.5)

$$\begin{aligned} & \int_0^t \left\| \frac{1}{\sqrt{n}} u'_n(t) \right\|^2 dt + \int_0^t \|B^{1/2} u'_n(t)\|^2 dt + \frac{1}{2} \varphi_n(u_n(t)) \\ & \leq c + \int_0^T \|f'(t)\| \left\| \frac{1}{n} A_n u_n(t) + J_n^A u_n(t) \right\| dt \\ & \leq c + \left( \frac{c}{n} + c \right) |f'|_T^2 + \frac{c}{n} \int_0^T \|A_n u_n(t)\|^2 dt + c \int_0^T \|J_n^A u_n(t)\|^2 dt \leq c. \end{aligned}$$

From this inequality and taking  $\varphi(J_n^A v) < \varphi_n(v)$  ( $v \in H$ ) into account, we see

$$(3.14) \quad \|J_n^A u_n(t)\| \leq c \quad \text{for every } t \in [0, T],$$

$$(3.15) \quad \left| \frac{1}{\sqrt{n}} u'_n \right|_T \leq c$$

and

$$(3.16) \quad |B^{1/2} u'_n|_T \leq c.$$

v) Taking the inner product of the both sides of (3.9) by  $t(u_n(t+h) - u_n(t))$ , ( $t > 0$ ) we have

$$\begin{aligned} & \frac{1}{2n} \frac{d}{dt} \|\sqrt{t}(u_n(t+h) - u_n(t))\|^2 - \frac{1}{2n} \|u_n(t+h) - u_n(t)\|^2 \\ & + \frac{1}{2n} \frac{d}{dt} \|\sqrt{t} B^{1/2}(u_n(t+h) - u_n(t))\|^2 - \frac{1}{2} \|B^{1/2}(u_n(t+h) - u_n(t))\|^2 \\ & + t(A_n u_n(t+h) - A_n u_n(t), u_n(t+h) - u_n(t)) \\ & = t(f(t+h) - f(t), u_n(t+h) - u_n(t)). \end{aligned}$$

Dividing the both sides of this equality by  $h^2$ , integrating it over  $(0, T)$  and letting  $h$  tends to zero, we see using (#), (A.2), (3.15) and (3.16) that

$$(3.17) \quad |\sqrt{t} u'_n|_T \leq c.$$

vi) From  $(E_n)$  and the estimates (3.11) and (3.15) we finally obtain

$$(3.18) \quad |A_n u_n|_T = \left| f - B u'_n - \frac{1}{n} u'_n \right|_T \leq c$$

and

$$(3.19) \quad |u_n|_T \leq c.$$

Here in (3.19) we have used (3.5), (3.18) and the identity (#).

From these estimates obtained in the steps i) ~ vi) it follows that a subsequence (denoted again by  $u_n$ ) can be extracted from  $\{u_n\}$  such that as  $n \rightarrow +\infty$ ,

$$\begin{aligned} u_n &\rightarrow u && \text{in } L^2(0, T; H) \text{ weakly,} \\ \frac{1}{n} u'_n &\rightarrow 0 && \text{in } L^2(0, T; H), \\ \sqrt{t} u'_n &\rightarrow \sqrt{t} u' && \text{in } L^2(0, T; H) \text{ weakly,} \\ B^{1/2} u_n &\rightarrow B^{1/2} u && \text{in } L^\infty(0, T; H) \text{ weakly star,} \\ B u_n &\rightarrow B u && \text{in } L^\infty(0, T; H) \text{ weakly star,} \\ B^{1/2} u'_n &\rightarrow (B^{1/2} u)' && \text{in } L^2(0, T; H) \text{ weakly,} \\ B u'_n &\rightarrow (B u)' && \text{in } L^\infty(0, T; H) \text{ weakly star,} \end{aligned}$$

and

$$A_n u_n \rightarrow \xi \quad \text{in } L^2(0, T; H) \text{ weakly.}$$

Note that  $(B u)'(t) = B u'(t)$  holds a.e.  $(0, T)$  by means of Remark 1.

Now passing to the limit in  $(E_n)$  as  $n \rightarrow +\infty$  we see

$$B \frac{du}{dt}(t) + \xi(t) = f(t) \quad \text{a.e. } (0, T).$$

Thus for concluding the proof, we need only to show that  $\xi(t) \in \partial\varphi(u(t))$  for a.a.  $t \in (0, T)$  and  $B^{1/2} u(t)$  satisfies (IC).

For every integer  $m, n > 0$  we have

$$\left( \frac{1}{m} u'_m - \frac{1}{n} u'_n, u_m - u_n \right) + \frac{1}{2} \frac{d}{dt} \|B^{1/2}(u_m - u_n)\|^2 + (A_m u_m - A_n u_n, u_m - u_n) = 0.$$

We see from (3.15) and (3.19) that

$$\left| \int_0^T \left( \frac{1}{m} u'_m - \frac{1}{n} u'_n, u_m - u_n \right) dt \right| \leq c \left( \frac{1}{\sqrt{m}} + \frac{1}{\sqrt{n}} \right)$$

and from (A.2) and (3.18) that

$$\begin{aligned} \int_0^t (A_m u_m - A_n u_n, u_m - u_n) dt &= \int_0^t \left( A_m u_m - A_n u_n, \frac{1}{m} A_m u_m - \frac{1}{n} A_n u_n \right) dt \\ &\quad + \int_0^t (A_m u_m - A_n u_n, J_m^A u_m - J_n^A u_n) dt \\ &\geq -c \left( \frac{1}{m} + \frac{1}{n} \right) + d \int_0^t \|J_m^A u_m - J_n^A u_n\|^2 dt. \end{aligned}$$

Hence combining them we have

$$\frac{1}{2} \|B^{1/2}(u_m(t) - u_n(t))\|^2 + d \int_0^t \|u_m(t) - u_n(t)\|^2 dt \leq c \left( \frac{1}{m} + \frac{1}{n} \right)$$

which follows

$$(3.20) \quad u_n \rightarrow u \quad \text{in } L^2(0, T; H) \text{ strongly}$$

and

$$(3.21) \quad B^{1/2}u_n \rightarrow B^{1/2}u \quad \text{in } C([0, T]; H) \text{ strongly.}$$

Then in virtue of (3.20) and demiclosedness of  $A$ , we see  $u(t) \in D(A)$  and  $\xi(t) \in Au(t)$  for a.a.  $t \in (0, T)$ . Finally  $B^{1/2}u(t)$  satisfies (IC) by means of (3.21). This completes the proof of Theorem 1.

#### 4. Application

We shall apply Theorem 1 to the initial-boundary value problem for some nonlinear partial differential equations.

Let  $\Omega$  be a bounded domain in  $\mathbf{R}^N$  with smooth boundary  $\partial\Omega$ . We shall denote  $\Delta$  the Laplacian in  $\mathbf{R}^N$  and  $\partial/\partial n$  the outward normal derivative at  $\partial\Omega$ .

*Example 1.* Consider an initial-boundary value problem

$$(*) \quad \begin{cases} -\Delta \frac{\partial u(x, t)}{\partial t} + (I - \Delta)u(x, t) = f(x, t) & \text{in } \Omega \times (0, T), \\ \frac{\partial u(x, t)}{\partial n} \in -\beta(u(x, t)), \quad \frac{\partial}{\partial n} \left( \frac{\partial u(x, t)}{\partial t} \right) = 0 & \text{on } \partial\Omega \times (0, T), \\ (-\Delta)^{1/2}u(x, t)|_{t=0} = (-\Delta)^{1/2}u_0(x) & \text{in } \Omega. \end{cases}$$

Here  $x \in \Omega, t \in (0, T), 0 < T < +\infty$  and  $\beta$  is an  $m$ -accretive operator in  $\mathbf{R}$  such that  $D(\beta)$  is dense in  $\mathbf{R}$ . (Then there exists a l.s.c. convex function  $j: \mathbf{R} \rightarrow (-\infty, +\infty]$  such that  $j \not\equiv +\infty$  and  $\partial j = \beta$ , [1].)

Let  $B = -\Delta_B = -\Delta$  with  $D(B) = \{u \in H^2(\Omega); (\partial u/\partial n)|_{\partial\Omega} = 0\}$ , then  $B$  is a nonnegative selfadjoint operator in  $H \equiv L^2(\Omega)$ . Let  $\varphi: L^2(\Omega) \rightarrow (-\infty, +\infty]$  be a function defined by

$$\varphi(u) = \begin{cases} \frac{1}{2} \int_{\Omega} (|u(x)|^2 + |\nabla u(x)|^2) dx + \int_{\partial\Omega} j(u) d\sigma & \text{if } u \in H^1(\Omega), j(u) \in L^1(\partial\Omega) \\ +\infty & \text{otherwise.} \end{cases}$$

Then it is well known [1] that  $\partial\varphi(u) = (I - \Delta)u, u \in D(\partial\varphi)$ , where  $D(\partial\varphi) = \{u \in H^2(\Omega); \partial u/\partial n \in -\beta(u) \text{ a.e. } \partial\Omega\}$ . Set  $A = \partial\varphi$ . We denote  $\Delta_A = I - \partial\varphi = I - A$ .

Then applying Theorem 1 to (\*) we have the following.

**Theorem 2.** Assume  $j(u) \geq 0$  a.e. on  $\partial\Omega$ . Then for every  $f \in W^{1,2}(0, T; L^2(\Omega))$  and  $u_0 \in D(A) \cap D(B)$  there exists a unique function  $u(x, t)$  which satisfies (\*) such that

$$u, (I - A_A)u \in L^2(0, T; L^2(\Omega)), \quad \sqrt{t} \frac{\partial u}{\partial t} \in L^2(0, T; L^2(\Omega)),$$

and

$$\begin{aligned} &(-A_B)^{1/2}u, \quad -A_B u \in AC([0, T]; L^2(\Omega)), \\ &\frac{\partial}{\partial t}((-A_B)^{1/2}u), \quad \frac{\partial}{\partial t}(-A_B u) \in L^2(0, T; L^2(\Omega)). \end{aligned}$$

*Proof.* (A.1), (A.2) and (A.3) are clear. Then in order to apply Theorem 1 we only show that (A.4) holds. For every  $u, v \in D(A) \cap D(B)$ , we have

$$\begin{aligned} &(Au - Av, B(u - v))_H \\ &= \int_{\Omega} \{(I - A_A)u(x) - (I - A_A)v(x)\} \{-A_B(u(x) - v(x))\} dx \\ &= \int_{\Omega} |\nabla(u(x) - v(x))|^2 dx + \int_{\Omega} A_A u(x) A_B u(x) dx + \int_{\Omega} A_A v(x) A_B v(x) dx \\ &\quad - \int_{\Omega} A_A u(x) A_B v(x) dx - \int_{\Omega} A_A v(x) A_B u(x) dx \\ &\geq \int_{\Omega} |\nabla(u(x) - v(x))|^2 dx \geq 0, \end{aligned}$$

which means (A.4) by taking Remark 2 into account.

*Example 2.* Let  $e(x)$  be an element of  $L^\infty(\Omega)$  such that  $e(x) \geq 0$  a.e. in  $\Omega$ . Consider an initial-boundary value problem

$$(**) \quad \begin{cases} e(x) \frac{\partial u(x, t)}{\partial t} + (I - A)u(x, t) = f(x, t) & \text{in } \Omega \times (0, T), \\ \frac{\partial u(x, t)}{\partial n} \in -\beta(u(x, t)) & \text{on } \partial\Omega \times (0, T), \\ \sqrt{e(x)}u(x, t)|_{t=0} = \sqrt{e(x)}u_0(x) & \text{in } \Omega. \end{cases}$$

Let  $Bu = e(x)u$  with  $D(B) = L^2(\Omega) \equiv H$  and let  $\varphi : L^2(\Omega) \rightarrow (-\infty, +\infty]$  be a function defined as in Example 1. Set  $A = \partial\varphi$ . Then  $B$  is a (bounded) non-negative selfadjoint operator in  $L^2(\Omega)$  and  $A = I - \mathcal{A}$  with  $D(A) = \{u \in H^2(\Omega);$

$\partial u/\partial n \in -\beta(u)$  a.e.  $\partial\Omega$  is an  $m$ -accretive operator in  $L^2(\Omega)$  which satisfies (A.1) and (A.2). Then Theorem 1 is applicable to (\*\*). Note that, in this case, the assumption (A.4) is not necessary because  $|B^{1/2}u'_n|_T \leq c$  implies  $|Bu'_n|_T \leq c$ .

### References

- [ 1 ] Barbu, V., *Nonlinear Semigroups and Differential Equations in Banach Spaces*, Noordhoff, Leyden, 1976.
- [ 2 ] Barenblatt, G. I., Zheltov, I. P. and Kochina, I. N., Basic concepts in the theory of seepage of homogeneous liquids in fissured rocks, *J. Appl. Math. Mech.*, **24** (1960), 1286–1303.
- [ 3 ] Benjamin, T. B., Bona, J. L. and Mahoney, J. J., Model equations for long waves in nonlinear dispersive systems, *Philos. Trans. Roy. Soc. London, Ser. A*, **272** (1972), 47–78.
- [ 4 ] Brézis, H., *Opérateurs maximaux monotones et semi-groupes de contraction dans les espace de Hilbert*, Math. Studies 5, North-Holland, Amsterdam, 1975.
- [ 5 ] Brill, H., A semilinear Sobolev equation in a Banach space, *J. Differential Equations*, **24** (1977), 412–425.
- [ 6 ] Carrol, R. W. and Showalter, R. E., *Singular and Degenerate Cauchy Problems*, Academic Press, New York/London, 1976.
- [ 7 ] Chen, P. J. and Gurtin, M. E., On a theory of heat conduction involving two temperatures, *Z. Angew. Math. Phys.*, **19** (1968), 614–627.
- [ 8 ] Kojo, T. (formerly Matahashi) and Tsutsumi, M., Initial value problem for an abstract differential equation in a Hilbert space, *Bull. Sci. Eng. Res. Lab., Waseda Univ.*, **87** (1979), 64–69.
- [ 9 ] Lagnese, J. E., General boundary value problems for differential equations of Sobolev type, *SIAM J. Math. Anal.*, **3** (1972), 105–119.
- [10] Lions, J. L., *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, Paris, 1969.
- [11] Showalter, R. E., Degenerate parabolic initial-boundary value problems, *J. Differential Equations*, **31** (1979), 296–312.
- [12] Sobolev, S. L., Some new problems in mathematical physics, *Izv. Akad. Nauk, SSSR*, **18** (1954), 3–50 (Russian).
- [13] Tsutsumi, M. and Kojo, T. (formerly Matahashi), On some nonlinear pseudo-parabolic equations, *J. Differential Equations*, **32** (1979), 65–75.
- [14] Tsutsumi, M. and Kojo, T. (formerly Matahashi), A certain class of singular nonlinear pseudo-parabolic equations, *Funkcialaj Ekvacioj*, **22** (1979), 313–325.
- [15] Zaidman, S., *Topics in abstract differential equations*, Pitman Research Notes in Math., 304, Longman, 1994.

nuna adreso:

Tomomi Kojo  
 Faculty of Systems Engineering  
 Shibaura Institute of Technology  
 Fukasaku, Omiya  
 Saitama 330  
 Japan

Masayoshi Tsutsumi  
Department of Applied Physics  
School of Science and Engineering  
Waseda University  
3-4-1 Okubo, Shinjuku-ku  
Tokyo 169  
Japan

(Ricevita la 26-an de februano, 1996)